

Western Australian Immunisation Schedule

Childhood

Age and eligibility	Antigen/Antibody	Brand(s)	Important information
Birth to <7 days	Nirsevimab*	Beyfortus®	For those residing or soon to reside in the Kimberley or Pilbara whose mother did not receive Abrysvo >2 weeks prior to birth. Give as close to birth as possible but can be given up to 8 months of age. See RSV decision aid and links below for further information.
	Hepatitis B	H-B-Vax® II Paed or Engerix® B Paed	
6-8 weeks	DTPa-hepB-IPV-Hib	Infanrix® hexa or Vaxelis®	
	20vPCV	Prevenar® 20	See here for childhood pneumococcal vaccination decision aid
Aboriginal* and/or MRC	Rotavirus	Rotarix®	Oral dose 1: Must be given by 14 weeks of age
	MenACWY	Nimenrix®	
Aboriginal and/or MRC	MenB	Bexsero®	Prophylactic paracetamol recommended
4 months	DTPa-hepB-IPV-Hib	Infanrix® hexa or Vaxelis®	
	20vPCV	Prevenar® 20	See here for childhood pneumococcal vaccination decision aid
Aboriginal* and/or MRC	Rotavirus	Rotarix®	Oral dose 2: Must be given by 24 weeks of age
	MenACWY	Nimenrix®	
Aboriginal and/or MRC	MenB	Bexsero®	Prophylactic paracetamol recommended
6 months	DTPa-hepB-IPV-Hib	Infanrix® hexa or Vaxelis®	
	20vPCV	Prevenar® 20	See here for childhood pneumococcal vaccination decision aid
MRC	MenB	Bexsero®	Prophylactic paracetamol recommended
	MenACWY	Nimenrix®	
12 months	MMR	Priorix® or M-M-R II®	The 2 brands of MMR vaccine are interchangeable
	20vPCV	Prevenar® 20	See here for childhood pneumococcal vaccination decision aid
Aboriginal and/or MRC	MenACWY	Nimenrix®	
	MenB	Bexsero®	Prophylactic paracetamol recommended
MRC	Hepatitis B	H-B-Vax® II Paed or Engerix® B Paed	<32 weeks gestation or <2000g birth weight
18 months	DTPa	Infanrix® or Tripacel®	
	Hib	ActHIB®	
Aboriginal	MMRV	Priorix-Tetra®	Do not administer MMRV as dose 1 of a measles-containing vaccine if <4 years of age
	Hepatitis A	Vaqta® Paed/Adol	Dose 1 of 2. Dose 2 to be administered at least 6 months after dose 1

<2 years	Catch-up for MenB is available for Aboriginal children <2 years of age		
4 years	DTPa-IPV	Quadracel® or Infanrix®IPV	
Aboriginal	Hepatitis A	Vaqta® Paed/Adol	Not required if previously received 2 doses (first dose at age \geq 12 months) at least 6 months apart
Aboriginal and/or MRC	20vPCV 23vPPV	Prevenar® 20 Pneumovax® 23	See here for childhood pneumococcal vaccination decision aid

Respiratory Syncytial Virus (RSV) – Kimberley and Pilbara regions ONLY

Age and eligibility	Antigen/Antibody	Brand(s)	Important information
See here for Kimberley and Pilbara RSV decision aid	Nirsevimab*	Beyfortus®	<p>The RSV decision aid contains guidance for residents, or soon-to-be residents, of the Kimberley or Pilbara, who meet one of the following criteria:</p> <ul style="list-style-type: none"> • Abrysvo for patients from 28 weeks gestation • Beyfortus for the following: <ul style="list-style-type: none"> - non-Aboriginal child without a medical-risk condition aged <8 months, or - Aboriginal child without a medical-risk condition up to 24 months of age, or - all children with a medical-risk condition up to 24 months of age. <p>Note: Children with medical-risk conditions may have up to 2 doses of Beyfortus until 24 months of age, at a minimum of 6 months apart.</p>

Age and eligibility	Antigen	Vaccine brand(s)	Important information
Year 7	HPV	Gardasil®9	Catch-up for adolescents <26 years of age
	dTpa	Boostrix® or Adacel®	
Year 10	MenACWY	Nimenrix® or MenQuadfi®	
≥18 years	Herpes-zoster	Shingrix®	2 doses, given 1–2 months apart
Pregnant	dTpa	Boostrix® or Adacel®	Each pregnancy, ideally between 20–32 weeks
	RSV	Abrysvo®	From 28 weeks gestation
	Influenza	Vaxigrip®Tetra, Flucelvax®Quad	Each pregnancy, any trimester
	Hepatitis B	Engerix® B	For non-immune persons. See the Australian Immunisation Handbook for recommendations and timing of doses
Adult refugees and humanitarian entrants	dTpa (Boostrix® or Adacel®), HepB (H-B-Vax® II or Engerix® B), IPV (IPOL®), MMR (MMR II® or Priorix®), VZV (Varivax®). HPV (Gardasil®9) <26 years of age		See the Australian Immunisation Handbook for timing of doses
≥50 years Aboriginal	13vPCV 23vPPV	Prevenar® 13 Pneumovax23®	See the Australian Immunisation Handbook for timing of doses
	Herpes-zoster	Shingrix®	2 doses, given 2–6 months apart
Born after 1965*	MMR	Priorix® or M-M-R II®	Give 1 or 2 doses as required. Whilst serology is not routinely recommended, see WA adult measles vaccination program for details. The 2 brands of MMR vaccine are interchangeable
≥65 years	Herpes-zoster	Shingrix®	2 doses, given 2–6 months apart
≥70 years	13vPCV	Prevenar® 13	

All children and adolescents aged over 12 months to under 18 years with **specified MRC** should receive 1 dose of Prevenar 20 at diagnosis. Refer to the [Australian Immunisation Handbook](#) for details

Additional *Haemophilus influenzae* type b (Hib), meningococcal and pneumococcal vaccines are recommended for people with specified risk conditions. Refer to the [Australian Immunisation Handbook](#) for details

Age and eligibility	Vaccine brand(s)	Important information
6 Months to <5 Years	Vaxigrip®Tetra, FluQuadri®, Flucelvax®Quad	
≥5 Years to Year 6*	FluQuadri®	
5 Years to <65 Years		
Aboriginal and/or MRC	Flucelvax®Quad, Vaxigrip®Tetra	For children <9 years, or persons following a stem cell or solid organ transplant, 2 doses are required 1 month apart in the first year they receive influenza vaccine. One dose annually in subsequent years. For information on influenza dosing see the Australian Immunisation Handbook or annual ATAGI advice on seasonal influenza vaccines.
Vulnerable cohorts* 12 to <65 Years	FluQuadri®	For non-NIP eligible residents in congregate living and people experiencing homelessness.
Pregnancy	Vaxigrip®Tetra, Flucelvax®Quad	Each pregnancy, any trimester
≥65 Years	Fluad® Quad, Flucelvax®Quad	Preferentially use adjuvanted QIV
WA Health staff**	Flucelvax®Quad, FluQuadri®, Fluad® Quad	For ≥65 years use Fluad® Quad

*Refers to WA State-funded Programs.

^Refers to all persons attending WA Health staff influenza vaccination services.

The term '**Aboriginal**' is inclusive of all Aboriginal and Torres Strait Islander people.

MRC refers to people with medical risk conditions. Refer to the current online edition of [Australian Immunisation Handbook](#) for all medical risk conditions, recommendations and timing of doses.

Persons aged <20 years can catch-up scheduled vaccines, refer to [Australian Immunisation Handbook](#) for timing of doses.

This document can be made available in alternative formats on request.

Produced by the Communicable Disease Control Directorate
© Department of Health 2026

Copyright to this material is vested in the State of Western Australia unless otherwise indicated. Apart from any fair dealing for the purposes of private study, research, criticism or review, as permitted under the provisions of the *Copyright Act 1968*, no part may be reproduced or re-used for any purposes whatsoever without written permission of the State of Western Australia.